


### AMENDMENT TO CLAIMS

This listing of claims replaces all prior listings, and versions, of claims in the application.

1. (Currently amended) ~~Use of a pharmaceutically effective amount of a polyphenol for the preparation of a medicament for the treatment of a cancerous lesion of the skin of a patient~~ A method for treating a cancerous lesion of the skin in a patient, said method comprising administering a pharmaceutically effective amount of a polyphenol to said patient.
2. (Currently amended) The ~~use~~ method according to claim 1, wherein the cancerous lesion is a pre-cancerous lesion.
3. (Currently amended) The ~~use~~ method according to claim 1, wherein the lesion of the skin is a non-virally induced lesion.
4. (Currently amended) The ~~use~~ method according to claim 3, wherein the lesion is a lesion not caused by papilloma virus.
5. (Currently amended) The ~~use~~ method according to claim 4, wherein the lesion is not a lesion selected from the group consisting of hyperplasia, Condyloma acuminata, warts and cervical intra-epithelial neoplasia.
6. (Currently amended) The ~~use~~ method according to claim 1, wherein the patient is a human.

7. (Currently amended) The ~~use~~ method according to claim 1, wherein the cancerous lesion of the skin is an advanced tumor.
8. (Currently amended) The ~~use~~ method according to claim 1, wherein the polyphenol is isolated from tea.
9. (Currently amended) The ~~use~~ method according to claim 1, wherein the polyphenol is extracted from a tea.
10. (Currently amended) The ~~use~~ method according to claim 9, wherein the tea is a green tea.
11. (Currently amended) The ~~use~~ method according to claim 1, wherein the polyphenol is isolated from a tea extract.
12. (Currently amended) The ~~use~~ method according to claim 1, wherein the polyphenol is selected from the group consisting of catechol, catechol gallate, epicatechol, epicatechol gallate, epigallocatechol, epigallocatechol gallate, gallocatechol and/or gallocatechol gallate.
13. (Currently amended) The ~~use~~ method according to claim 1, wherein the polyphenol has the general formula (I)




(I)

in which

$R_1$ ,  $R_2$  and  $R_6$  are independently from each other -H or -OH,

$R_3$  is -H or =O and

$R_4$  is independently from each other -H, -OH or a group of the formula (III)




(III)

$R_5$  and  $R_7$  are independently from each other -H, -OH or -OCH<sub>3</sub>, and

----- optionally represents a bond

14. (Currently amended) The ~~use~~ method according to claim 12, wherein the catechol has the general formula (II)




(II),

in which

$R_8$  is -H or -OH, and

$R_9$  is -H or a group of the formula (III)



(III)

15. (Currently amended) The use method according to claim 12, wherein the catechol is selected from the group consisting of of (+)-catechol, (-)-catechol, (+)-catechol gallate, (-)-catechol gallate, (+)-epicatechol, (-)-epicatechol, (+)-epicatechol gallate, (-)-epicatechol gallate, (+)-epigallocatechol, (-)-epigallocatechol, (+)-epigallocatechol gallate, (-)-epigallocatechol gallate, (+)-gallocatechol, (-)-gallocatechol, (+)-gallocatechol gallate and/or (-)-gallocatechol gallate.

16. (Currently amended) The use method according to claim 1, wherein the polyphenol is present in the form of a mixture of polyphenols, in particular catechols, especially containing catechol, catechol gallate, epicatechol, epicatechol gallate, epigallocatechol,

epigallocatechol gallate, gallocatechol and/or gallocatechol gallate, whereas the gallates of catechol, epicatechol, epigallocatechol or of gallocatechol are most preferred in the mixture of polyphenols.

17. (Currently amended) The use method according to claim 16, wherein the mixture of polyphenols is a tea extract.

18. (Currently amended) The use method according to claim 17, wherein the tea extract is a green tea extract.

19. (Currently amended) The use method according to claim 17, wherein the mixture of polyphenols is selected from the group consisting of catechol, catechol gallate, epicatechol, epicatechol gallate, epigallocatechol, epigallocatechol gallate, gallocatechol and/or gallocatechol gallate.

20. (Currently amended) The use method according to claim 19, wherein the catechols are selected from the group consisting of (-)-epicatechol, (-)-epicatechol gallate, (-)-epigallocatechol, (-)-epigallocatechol gallate, (+)-gallocatechol and/or (-)-gallocatechol gallate.

21. (Currently amended) The use method according to claim 19, wherein the catechols are selected from the group consisting of about 2-20% (w/w) epicatechol, about 2-20% (w/w) epicatechol gallate, about 1-25% (w/w) epigallocatechol, about 40-75% (w/w) epigallocatechol gallate, about 0.05-5% (w/w) gallocatechol and/or about 0.5-20% (w/w) gallocatechol gallate.

22. (Currently amended) The ~~use~~ method according to claim 19, wherein the catechols are selected from the group consisting of about 10.8% (w/w) of epicatechol, about 6.5% (w/w) of epicatechol gallate, about 9.2% (w/w) of epigallocatechol, about 54.8% (w/w) of epigallocatechol gallate and/or about 4.0% (w/w) of gallocatechol gallate.

23. (Currently amended) The ~~use~~ method according to claim 19, wherein the mixture contains about 10.8% (w/w) of (-)-epicatechol, about 6.5% (w/w) of (-)-epicatechol gallate, about 9.2% (w/w) of (-)-epigallocatechol, about 54.8% (w/w) of (-)-epigallocatechol gallate and about 4.0% (w/w) of (-)-gallocatechol gallate.

24. (Currently amended) The ~~use~~ method according to claim 19, wherein the catechols are selected from the group consisting of about 2-12% (w/w) epicatechol, about 4-15% (w/w) epicatechol gallate, about 1-8% (w/w) epigallocatechol, about 60-68% (w/w) epigallocatechol gallate, about 0.05-1% (w/w) gallocatechol and/or about 1-7% (w/w) gallocatechol gallate.

25. (Currently amended) The ~~use~~ method according to claim 19, wherein the catechols are selected from the group consisting of about 5-8% (w/w) epicatechol, about 5-7% (w/w) epicatechol gallate, about 2-3% (w/w) epigallocatechol, about 61-65% (w/w) epigallocatechol gallate and/or about 2-4% (w/w) of gallocatechol gallate.

26. (Currently amended) The ~~use~~ method according to claim 19, wherein the catechols are selected from the group consisting of about 5-8% (w/w) epicatechol, about 5-6% (w/w) epicatechol gallate, about 6-8% (w/w) epigallocatechol, about 61-65% (w/w) epigallocatechol gallate and/or about 2-4% (w/w) of gallocatechol gallate.

27. (Currently amended) The ~~use~~ method according to claim 1, wherein the polyphenol is combined with an additive.

28. (Currently amended) The ~~use~~ method according to claim 27, wherein the additive is selected from the group consisting of petroleum jelly, wax, oleyl alcohol, propylene glycol monostearate, propylene glycol monopalmitostearate and/or isopropyl myristate.

29. (Currently amended) The ~~use~~ method according to claim 1, wherein the polyphenol is administered topically.

30. (Currently amended) The ~~use~~ method according to claim 1, wherein the polyphenol is contained in a carrier selected from the group consisting of an emulsion, a gel, a cream and/or an ointment.

31. (Currently amended) The ~~use~~ method according to claim 1, wherein the method for treating said lesions is combined with a different anticancer treatment.

32. (Currently amended) The ~~use~~ method according to claim 31, wherein the different anticancer treatment is selected from the group consisting of surgery, electrodesiccation, curettage, excision, Mohs micrographic surgery, radiation, proton therapy, chemotherapy, photodynamic therapy, cryosurgery, laser, immunotherapy, vaccine therapy and/or biologic therapy.

33. (Currently amended) The ~~use~~ method according to claim 32, wherein the chemotherapy is carried out with an agent selected from the group consisting of podophyllin, 5-fluorouracil, bleomycin, interferon, imiquimod, and mixtures thereof.

34. (Currently amended) The ~~use~~ method according to claim 32, wherein the radiation is selected from the group consisting of X-ray radiation and  $\gamma$ -radiation.

35. (Currently amended) The ~~use~~ method according to claim 1 or 2, wherein the lesion of the skin is selected from the group consisting of skin cancer, cutaneous carcinoma, basal cell carcinoma, squamous cell carcinoma, actinic keratosis, solar keratosis, epithelioma, epithelial tumors, skin neoplasm, Bowen's disease, acanthoma, cancrroid, cutaneous horn, hyperkeratosis, keratosis, molluscum contagiosum, lid tumors, xanthelasma, xanthoma, fibroma, verucca senilis, seborrheic keratosis, cheilocarcinoma, papillomatosis, penis carcinoma, radiodermatitis, sailor's skin, tar cancer, vaginal carcinoma, vulvar cancer, erythroplasia queyrat and/or carcinoma of the tongue, in particular actinic or solar keratosis and/or basal cell carcinoma.